

Unrealistic Nonphysiological Amounts of Reagents and a Disregard for Published Literature

Isaac Ginsburg

Institute of Dental Sciences, The Hebrew University, Hadash Faculty of Dental Medicine, Ein Kerem Campus, Jerusalem, Israel

ere are some comments and useful suggestions after reading an article in *mBio* by Brown et al. entitled "Mechanisms underlying the exquisite sensitivity of Candida albicans to combinatorial cationic and oxidative stress that enhances the potent fungicidal activity of phagocytes" (1).

In this paper, we are informed that a simultaneous exposure to $5 \text{ mM H}_2\text{O}_2$ and to cationic NaCl at 1 M is much more potent than the individual stresses themselves and that this combinatorial stress kills C. albicans synergistically in vitro. Such combinations are obviously absolutely unrealistic and not physiological. As a comparison I wonder why the authors had not also tested naturally occurring antimicrobial cationic peptides such as LL37 found in large amounts in neutrophil granules? Had the authors read the classical papers describing the possible mechanisms of bactericidal effects of neutrophils, they would have realized that there is actually no free-floating H₂O₂ in phagosomes following phagocytosis. This is because activation of NADPH-oxidase yields superoxide, which very rapidly interacts with myeloperoxidase (MPO) and with a halide (Cl⁻) to generate microbicidal amounts of hypochlorous acid (HOCl) (2-5)! Therefore, HOCl should have definitely been considered and tested in the system described by the authors. Also, the term flux used may be inappropriate since, in their study, both H₂O₂ and NaCl were actually applied as a bolus. Fluxes of oxidants are generated mainly by activated neutrophils and macrophages and by xanthine and xanthine oxidase in endothelial cells (2) Also, I wonder whether Na used is specific and whether potassium ions can also have the same effects in their system? The authors also claimed that catalase-derived peroxide detoxification, which is inhibited by cations, leads to intracellular ROS accumulation because catalase activity had been affected. If so, why had the catalase inhibitor azide or aminotriazole not been tested? In their study, the authors grew Candida cells in Trisbuffered yeast extract-peptone-dextrose medium (YPDT; pH 7.4). However, the authors have not cited key papers showing that D-glucose, in media on which candida grow, may also suppress catalase formation (6, 7).

Using unrealistic, nonphysiological amounts of reagents will not increase our understanding of how biological processes really occur in vivo, despite the need to employ in vitro models. Also, disregarding key published data on neutrophil functions and Candida biology is unacceptable. Can this be a "menace to the future of honest science" (8) and also a "transgression" (9)? See also a recent publication by Casadevall and Fang (10).

REFERENCES

- 1. Kaloriti D, Jacobsen M, Yin Z, Patterson M, Tillmann A, Smith DA, Cook E, You T, Grimm MJ, Bohovych I, Grebogi C, Segal BH, Gow NA, Haynes K, Quinn J, Brown AJ. 2014. Mechanisms underlying the exquisite sensitivity of Candida albicans to combinatorial cationic and oxidative stress that enhances the potent fungicidal activity of phagocytes. mBio 5(4):e01334-14. http://dx.doi.org/10.1128/mBio.01334-14.
- 2. Hampton MB, Kettle AJ, Winterbourn CC. 1998. Inside the neutrophil phagosome: oxidants, myeloperoxidase, and bacterial killing. Blood 92:
- 3. Segal AW. 2005. How neutrophils kill microbes. Annu Rev Immunol 23: 197-223. http://dx.doi.org/10.1146/annurev.immunol.23.021704.115653.
- 4. Wellington M, Dolan K, Krysan DJ. 2009. Live Candida albicans suppresses production of reactive oxygen species in phagocytes. Infect Immun 77:405–413. http://dx.doi.org/10.1128/IAI.00860-08.
- 5. Klebanoff SJ, Kettle AJ, Rosen H, Winterbourn CC, Nauseef WM. 2013. Myeloperoxidase: a front-line defender against phagocytosed microorganisms. J Leukoc Biol 93:185-195. http://dx.doi.org/10.1189/
- 6. Rodaki A, Bohovych IM, Enjalbert B, Young T, Odds FC, Gow NA, Brown AJ. 2009. Glucose promotes stress resistance in the fungal pathogen Candida albicans. Mol Biol Cell 20:4845-4855. http://dx.doi.org/ 10.1091/mbc.E09-01-0002.
- 7. Nakagawa Y, Koide K, Watanabe K, Morita Y, Mizuguchi I, Akashi T. 1999. The expression of the pathogenic yeast Candida albicans catalase gene in response to hydrogen peroxide. Microbiol Immunol 43:645-651. http://dx.doi.org/10.1111/j.1348-0421.1999.tb02452.x.
- 8. Ginsburg I. 2001. The disregard syndrome, a menace to the future of honest science. Scientist 15:51.
- Garfield E. 1991. Bibliographic negligence: a serious transgression. Scientist 5:14.
- 10. Casadevall A, Fang FC. 2014. Diseased science. Microbe 9:390-392.

Published 21 April 2015

Citation Ginsburg I. 2015. Unrealistic nonphysiological amounts of reagents and a disregard for published literature. mBio 6(2):e00360-15. doi:10.1128/mBio.00360-15.

Copyright © 2015 Ginsburg. This is an open-access article distributed under the terms of the Creative Commons Attribution-Noncommercial-ShareAlike 3.0 Unported license, which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

Address correspondence to Ginsburg@mail.huji.ac.il.